



The Importance of Prenatal Stress on the Development of ADHD in Children Literature Review 2019

Dr. Tamar Kacharava^{a*}, Ketevan Nemsadze^b

^a*PhD Student, G. Chubinashvili street 15, Tbilisi 0164, Georgia*

^b*MD, Full Prof. N. Nikoladze street 3, Tbilisi 0108, Georgia*

^a*Email: tako_kat@yahoo.de*

^b*Email: ketinemsadze@gmail.com*

Abstract

Attention deficit hyperactivity disorder (ADHD) is a neurodevelopmental disorder that affects children and often persists into adulthood. Research suggests that there is a link between maternal stress levels and the development of this condition. According to the Centers for Disease Control and Prevention (CDC), 6.1 million children living in the United States in 2016 had received a diagnosis of ADHD. The symptoms of ADHD can vary from person to person, but they typically include difficulties with concentrating, paying attention, and controlling impulses. It is not clear what causes ADHD, but scientists believe that genetics, certain environmental factors, and brain changes may play a role in its development. Researchers have also investigated the role of nutrients, such as vitamin D and hormone, such as prenatal testosterone. In this article, we will review the latest literature assessing the relationship between prenatal exposure to psychosocial stress during pregnancy and to the risk of developing behavioral problems related to attention deficit hyperactivity disorder (ADHD) in Childhood.

Keywords: Prenatal stress; Attention deficit hyperactivity disorder (ADHD); Vitamin D; Prenatal testosterone; Etiological factors.

* Corresponding author.

1. Introduction

Attention deficit hyperactivity disorder (ADHD) is one of the most common behavioral conditions, particularly among children [1]. According to World Health Organization (WHO) surveys, its prevalence averages 2.2% among children worldwide and is followed by adult ADHD in 57% of the cases, although prevalence of adult ADHD is likely to be underestimated [2]. ADHD typically starts in childhood and often persists throughout life [3]. The symptoms are inattention, hyperactivity, and impulsivity [4]. People with ADHD might reach lower education levels than healthy children and their occupational attainment and social level might be lower in adulthood, and this might have an economic burden to societies [5]. The developmental aspect of ADHD must be taken into account when characterizing clinical presentation. The validity of ADHD among preschoolers has been an area of particular controversy in the literature. Although there is increasing evidence that ADHD constitutes a valid diagnosis even before the age of 6, there are several challenges in making a diagnosis during this developmental period. For example, the difficulties associated with making observations across multiple settings for children not attending preschool – and subsequent lack of information about pervasiveness. In addition, hyperactivity and impulsivity are much more prominent at this stage and inattention might not be so evident due to there being less environmental demands on the child. Thus, it is not surprising that ADHD predominantly hyperactive/impulsive presentation is the most frequent presentation in preschoolers. Several studies have however shown that currently available criteria reliably identify ADHD in children as young as 3 years old and that these individuals have clinically significant impairment across all relationships and settings [6]. Years ago, ADHD was seen as a problem occurring only in little boys who were hyperactive. It is now clear that ADHD occurs in many individuals who are not hyperactive. Although it is more often recognized in males, it also is found in a significant number of girls and women. High intelligence is not a protection against ADHD. Many with ADHD are very bright, but still struggle a lot in exercising executive functions described above which are essential for success in school, work and many activities of daily life. A very large percentage of children, teens and adults with ADHD have one or more additional problems such as anxiety, depression, sleep difficulties, substance use disorders, obsessive compulsive disorder, autism spectrum disorders, and/or specific learning disorders in reading, math or written expression. One of these other problems may be identified first, possibly overlooking the underlying ADHD. Or the ADHD may be recognized while another underlying disorder is not noticed or treated [7]. The causes of ADHD are not fully understood; however, several environmental (e.g., exposure to maternal stress during pregnancy, certain foods or inhalants) and genetic risk factors have been proposed [8]. Modern obstetrics has seen dramatic changes in the past decades. The fetus has become a patient and modern obstetrics has matured into feto-maternal medicine. The latest development is the advent of the new concept of „fetal programming” which is based on the understanding that the intrauterine environment can impose postgenomic changes that can have far reaching consequences on health and wellbeing which can affect the individual in adulthood as well as subsequent generations. Understanding the concept of fetal programming requires the understanding of epigenetics, which in essence is the nonmutational manipulation of genes without affecting their basic structure. Fetal programming may create predisposition factors for adult disease and is also crucial for behavioral and cognitive normalcy or deviation thereof. The importance of fetal programming cannot be overstated. It is now common understanding that prenatal life is no safe haven for the fetus and that the environment in which the pregnant mother lives has a direct impact on the

development of the fetus. In effect, there is no other time throughout the life span of an individual where it is so intimately exposed to the environment. Whatever affects the pregnant mother may well affect her growing embryo and fetus, in many cases in a greatly amplified manner. The impact of exogenous toxins on the developing fetus is dependent on qualitative and quantitative factors and also on when they occur during the development of the fetus. First trimester exposure will generally have teratogenic effects while second and third trimester exposure will more often be expressed in growth restriction and organ failure [9]. The importance of development during the fetal period is well established with regards to the association between the baby's growth in the womb, and later vulnerability to physical disorders such as cardiovascular disease and other aspects of metabolic syndrome [10]. It is now clear that environmental effects on fetal development are important with respect to emotional, behavioural and cognitive outcomes too. Many groups around the world are studying how the emotional state of the mother during pregnancy can have long-lasting effects on the psychological development of her child [11-12]. Some are using large population cohorts, which have the advantage of being able to statistically allow for many confounding factors including postnatal maternal mood. [13] Others are smaller observational studies which can examine the child in more detail [14]. Stress is a generic term which includes anxiety and depression, but also includes distress due to poor relationships or the response to an acute disaster. All these have been shown to be associated with altered outcome for the child. Many independent prospective studies have now shown that if a mother is stressed, anxious or depressed while pregnant, her child is at increased risk for having a range of problems, including emotional problems, ADHD, conduct disorder and impaired cognitive development. Both altered brain structure [15] and function [16] have been shown to be associated with prenatal stress, and also the mother's experience of early childhood trauma [17]. While genetic transmission and the quality of postnatal care are likely to contribute to some of these findings of association, there is good evidence that there is a causal influence of the mother's emotional state while pregnant also. Some studies have found stronger associations with prenatal maternal mood than paternal [18]. Several large cohort studies have found associations independent of possible confounding factors, such as birthweight, gestational age, maternal education, smoking, alcohol consumption, and most importantly, postnatal anxiety and depression [13]. Thus, although the mother's postnatal emotional state and the quality of early postnatal care are clearly important for many of these outcomes, the evidence suggests that there are substantial prenatal effects also. Within a normal population, the children of the most anxious mothers during pregnancy (top 15%), had double the risk of emotional or behavioural problems, compared with the children of the less anxious mothers [13]. Most children were not affected, and those that were, were affected in different ways. However a doubling of risk is of considerable clinical significance. Several studies are finding that boys and girls can be affected in different ways [11]. There are gene environment interactions too, in that a child with a specific genetic vulnerability is more likely to be affected in a particular way [19]. It is clear that it is not just toxic or extreme prenatal stress that are important, as several studies have shown that problems such as daily hassles, pregnancy specific anxiety or relationship strain [14] can have an adverse effect on the developing fetus. Effects of acute disasters such as [16-18,20] have also been demonstrated. Different studies have shown different gestational ages of vulnerability. This may vary for different outcomes. Increased vulnerability to schizophrenia has been found to be associated with extreme stress in the first trimester [21]. The risk for other outcomes, such as ADHD, has been found to be associated with stress later in pregnancy [13,22]. When discussing causation, studies of DNA risk variants have a clear advantage over studies of the environment. Our

genome comes into existence prior to our birth. So, when scientists discover an association between ADHD and a DNA variant, it is clear that having ADHD cannot “cause” one to have a specific DNA variant but that having a DNA variant could logically increase risk for ADHD. Studies of the environment are less clear-cut. For example, if a study documents that poverty is associated with ADHD, that could mean that poor nutrition, stress and other concomitants of poverty increase the risk for ADHD. But it is also possible that having ADHD leads to lower levels of education, poorer job performance and thereby increases the risk for parents with high genetic risk for ADHD to live in poverty. Thus, one must always keep in mind the potential for such ‘reverse causation’ when evaluating environmental risk factors and evaluate whether these have been considered by the relevant studies. That said, keep in mind that when one member of an identical twin pair has ADHD, the risk to the co-twin is only about 50%. Thus, environmental risk factor must contribute to the etiology of ADHD [7]. Some environmental risks are due to exposures to toxins, lack of nutrients or trauma. Many studies have examined the effects of iron and zinc on ADHD because both of these elements are essential for producing norepinephrine and dopamine in the brain. In a meta-analysis, Scientist [23] reported that measures of iron deficiency were associated with ADHD. They also found that ADHD was associated with low levels of zinc in the blood. Among the many toxins studied in ADHD patients, the strongest evidence implicates lead contamination. In their meta-analysis, they found that compared with controls, ADHD cases were more likely to have been exposed to lead. Many studies have tested the idea that pregnancy and delivery complications (PDCs) might cause ADHD by harming the brain at early stages of its development. Although the literature presents conflicting results, it tends to support the idea that PDCs are risk factors for ADHD. When PDCs have been implicated in ADHD they typically lead to oxygen deprivation and tend to involve *chronic* exposures to the fetus, not *acute* events. Among the most investigated PDCs, prematurity and low birth-weight are the most studied. A recent meta-analysis of the literature on the association between both very-premature and/or very low-weight babies and ADHD showed a 3 times increased risk for those infants to have ADHD in the future [24]. However, it is important to note that prematurity and low-birth weight are risk factors to other mental disorders. Anyhow, whenever assessing very premature and/or very low birth weight children, clinicians might consider assessing for ADHD. Maternal smoking during pregnancy has been widely studied as a risk factor for ADHD. It is well documented that smoking during pregnancy places the fetus at risk for birth complications, including low birth weight, which has been associated with ADHD. Maternal smoking also places the fetus at risk for a hypoxia, which has been associated with ADHD. Although one group of scientist’s [25] meta-analysis concluded that children whose mothers’ smoked during pregnancy had a 2.4 fold increased risk for ADHD, this is still an area of debate because ADHD and its polygenic risk are known to be associated with smoking behaviors. Thus, mother with ADHD might smoke more than mothers without ADHD and the risk is associated to genetic factors related to ADHD and not to smoking. Those who favor the maternal smoking hypothesis point out that it is a plausible risk factor because nicotine regulates the activity of the dopamine transporter, the site of action of the stimulant drugs that treat ADHD. People who experience mild traumatic brain injuries (mTBIs) are at risk for developing ADHD. This was the conclusion of a meta-analysis which showed that mTBI associated with ADHD. [26] Another well-documented environmental risk factor is severe institutional deprivation in early childhood. We know this from studies of children who spent the early years of life in Romanian orphanages that offered poor nutrition and nearly no human contact. Many of these children developed ADHD later in life [27]. Environmental risk factors for ADHD that have been confirmed by meta-analyses include: preterm birth [28], prenatal exposure to maternal

smoking [29] prenatal methylmercury exposure from maternal fish consumption [30] exposure to lead [31] and perinatal vitamin D deficiency [32]. From meta-analyses, we can also exclude some environmental factors as increasing the risk for ADHD. These include: sugar consumption [33], methylmercury in vaccines [30], maternal thyroid hormone insufficiency [34], sleep restriction [35], cesarean section [36] and solar intensity [37]. It is easy to see how toxic exposures, pregnancy and delivery complications, traumatic brain injuries and severe institutional deprivation could affect the developing brain and increase risk for ADHD. In addition to such biological adversity, studies have also implicated adverse psychosocial experiences as risk factors for the disorder. Examples of psychosocial stressors that affect children are marital distress, family dysfunction and low social class. Other potential risk factors for ADHD are low maternal education, low social class, and single parenthood. Several studies show that the mothers of ADHD children have more negative communication patterns, more conflict with their child and a greater intensity of anger than do control mothers and that families of ADHD children are more likely to have higher levels of chronic conflict, decreased family cohesion, and exposure to parental psychopathology. However, most of the environmental factors might act more as unspecific triggers for mental health problems in general than specific environmental risk factors for ADHD. An association between vitamin D and attention deficit hyperactivity disorder (ADHD) has been proposed by several researchers in recent years; however, the investigations have led to inconsistent results. It has been suggested that dietary interventions such as ω -3 FA [38] and vitamin and mineral supplementation [39], restriction diets, and the avoidance of synthetic food color additives [40] might affect ADHD symptoms [41]. Recently a number of studies have proposed that vitamin D might play a role in ADHD pathogenesis. The mechanisms by which vitamin D might affect a number of neurological diseases, including ADHD, are not clear. Nevertheless, there is evidence demonstrating the widespread presence of vitamin D receptors and 1α -hydroxylase (the enzyme responsible for the formation of the active vitamin) in the human brain; therefore, it is suggested that vitamin D might have neurohormonal properties in the human brain [42]. Furthermore, a recent study proposed that vitamin D directly upregulates expression of tyrosine hydroxylase (a rate-limiting enzyme in dopamine synthesis) by binding to the nuclear vitamin D receptor [43]. It is also suggested that this vitamin is involved in the synthesis of serotonin in the brain [44]. Several observational studies have examined the association between vitamin D status and ADHD in children and adolescents in recent years; however, they have achieved inconsistent results [45-48]. The majority of case-control studies have revealed that serum vitamin D concentrations are lower in ADHD children than in healthy controls [46,47, 49-51]. However, a study done by other scientists [45] could not show the same association. Moreover, in a cohort study, researchers [48] reported that there is no difference in cord blood vitamin D concentrations at birth between children with ADHD and healthy controls. In contrast, another cohort study conducted by another group of researchers [52] revealed that higher maternal circulating concentrations of 25-hydroxyvitamin D [25(OH)D] in pregnancy might be associated with a lower risk of developing ADHD-like symptoms in offspring. Although the findings of a connection between vitamin D status and ADHD have been controversial [53]. On the basis of observations of sex differences, many researchers have raised questions with respect to the potential roles of the neuroendocrine system in the etiology of ADHD [54-56]. The male gonad commences testosterone secretion as early as the 7th week of pregnancy. Testosterone levels peak around 14-18 weeks of gestation and testosterone levels in the developing male fetus may become close to adult levels. Testosterone exerts manifold effects on the developing organism, like sex-specific development of genitalia and growth rate. The effect of fetal sex steroids on

mammalian brain development is most striking. Dependent on the availability of testosterone receptors, testosterone affects cell apoptosis and is involved in the establishment of neural connectivity. The results of this development are far reaching and are of crucial importance for „male” behavior after birth and throughout life, including childhood play behavior, and may also be involved in the establishment of sexual orientation and sexual identity. High exogenous or endogenous testosterone levels in female embryos (or neonates) will lead to behavior patterns typical of males [57]. High levels of prenatal T may affect the dopaminergic neural circuitry and decelerate neural development globally [58]. Excessive prenatal exposure to testosterone has been implicated in the development of dyslexia and autism [59]. On the other hand, reduced levels of fetal testosterone can be found in situations where the pregnant woman is under acute and chronic stress, such as in times of war or when exposed to natural or personal disasters. Prenatal maternal stress and anxiety have been shown to be associated with impaired neurodevelopmental outcomes, including attention deficit and hyperactivity in boys [60]. Several studies support the hypothesis that prenatal T exposure contributes to the development of ADHD in children. However, some researchers have not found associations between prenatal testosterone level and ADHD features or externalizing behavior problems in children [61-63]. In sum, whether prenatal T exposure plays a role in the etiology of ADHD is still open to debate. Individuals with ADHD have at least 1.5 times the average risk of developing dependence on nicotine, alcohol, marijuana, cocaine, and other drugs [64], and tend to have earlier onset, faster progression, and less remittance of substance use disorders (SUDs) [65-66]. They also have higher rates of obesity [67] and three times higher rates of binge eating than individuals without ADHD [68]. Furthermore, ADHD is associated with unsafe sexual behavior, including earlier sexual activity and more partners [69-70]. ADHD youth are also less likely to use contraception, leading to higher rates of sexually transmitted infections (STIs) and unintended pregnancies [70]. SUDs, obesity, and STIs are directly associated with chronic health problems as well as morbidity and mortality. SUDs also increase rates of violence, accidents, and unprotected sex. These difficulties, along with unintended pregnancies, additionally reduce socioeconomic attainment and quality of life. In light of the high prevalence of ADHD and associated risks, health promotion efforts for this population are much needed and have the potential for substantial public health impact. Future research questions and implications for clinicians are - when and how medical practitioners may optimally intervene to avoid the development of ADHD in children and how improve child health through early screenings, increasing medication adherence, and treating psychosocial impairments.

References

- [1]. Subcommittee on Attention-Deficit/Hyperactivity Disorder, Steering Committee on Quality Improvement and Management, Wolraich M, Brown L, Brown RT, DuPaul G, Earls M, Feldman HM, Ganiats TG, et al. ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics* 2011;128(5):1007–22. doi: 10.1542/peds.2011-2654.
- [2]. Fayyad, J. et al. The descriptive epidemiology of DSM-IV Adult ADHD in the World Health Organization World Mental Health Surveys. *ADHD Atten. Deficit Hyperact. Disord.*, <https://doi.org/10.1007/s12402-016-0208-3> (2016)
- [3]. Swanson J, Arnold LE, Kraemer H, Hechtman L, Molina B, Hinshaw S, Vitiello B, Jensen P, Steinhoff K, Lerner M. Evidence, interpretation, and qualification from multiple reports of long-term outcomes

- in the Multimodal Treatment Study of Children with ADHD (MTA) Part II: Supporting details. *J AttenDisord* 2008;12(1): 15–43.
- [4]. ATTENTION-DEFICIT SO. ADHD: clinical practice guideline for the diagnosis, evaluation, and treatment of attention-deficit/hyperactivity disorder in children and adolescents. *Pediatrics* 2011: peds. 2011–654.
- [5]. Klein RG, Mannuzza S, Olazagasti MAR, Roizen E, Hutchison JA, Lashua EC, Castellanos FX. Clinical and functional outcome of childhood attention-deficit/hyperactivity disorder 33 years later. *Arch Gen Psychiat* 2012;69(12):1295–303.
- [6]. Dias TG, Kieling C, Graeff-Martins AS, Moriyama TS, Rohde LA, Polanczyk GV. Developments and challenges in the diagnosis and treatment of ADHD. *Braz J Psychiatr*. 2013;35Suppl 1:S40-50.
- [7]. Rohde, Buitelaar, Gerlach and Faraone, World Federation of ADHD. 2019
- [8]. Biederman J. Attention-deficit/hyperactivity disorder: a selective overview. *Biol Psychiatry* 2005;57(11):1215–20.
- [9]. Schneck-Gustaffson K, DeCola PR, Pfaff DW, Pisetsky DS (eds): *Handbook of Clinical Gender Medicine*. Basel, Karger, 2012, pp 37-51 „Fetal Programming”
- [10]. Barker DJ. The developmental origins of adult disease. *European Journal of Epidemiology* 2003;18(8):733-6.
- [11]. Van den Bergh BRH, van den Heuvel MI, Lahti M, Braeken M, de Rooij SR, Entringer S, Hoyer D, Roseboom T, Räikkönen K, King S, Schwab M. Prenatal developmental origins of behavior and mental health: the influence of maternal stress in pregnancy. . 2017. doi: 10.1016/j.neubiorev.2017.07.003. [Epub ahead of print] *Neuroscience and biobehavioral reviews*
- [12]. Monk C, Lugo-Candelas C, Trumpff C. Prenatal developmental origins of future psychopathology: mechanisms and pathways. . 2019. doi:10.1146/annurev-clinpsy-050718-095539. [Epub ahead of print] *Annual Review of Clinical Psychology*
- [13]. O'Donnell KJ, Glover V, Barker ED, O'Connor TG. The persisting effect of maternal mood in pregnancy on childhood psychopathology. *Developmental Psychopathology*. 2014;26(2):393-403.
- [14]. Bergman K, Sarkar P, Glover V, O'Connor TG. Maternal stress during pregnancy predicts cognitive ability and fearfulness in infancy. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2007;46(11):1454-1463.
- [15]. Buss C, Davis EP, Muftuler LT, Head K, Sandman CA. High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6-9-year-old children. *Psychoneuroendocrinology*. 2010;35(1):141-53.
- [16]. Mennes M, Van den Bergh B, Lagae L, Stiers P. Developmental brain alterations in 17 year old boys are related to antenatal maternal anxiety. *Clinical Neurophysiology*. 2009;120(6):1116-1122.
- [17]. Moog NK, Entringer S, Rasmussen JM, Styner M, Gilmore JH, Kathmann N, Heim CM, Wadhwa PD, Buss C. Intergenerational effect of maternal exposure to childhood maltreatment on newborn brain anatomy. *Biological Psychiatry*. 2018;83(2):120-127.
- [18]. Capron LE, Glover V, Pearson RM, Evans J, O'Connor TG, Stein A, Murphy SE, Ramchandani PG. Associations of maternal and paternal antenatal mood with offspring anxiety disorder at age 18 years. *Journal of Affective Disorders*. 2015;187:20-26.

- [19]. O'Donnell KJ, Glover V, Lahti J, Lahti M, Edgar RD, Räikkönen K, O'Connor TG. Maternal prenatal anxiety and child COMT genotype predict working memory and symptoms of ADHD. *PLoS One*. 2017;12(6):e0177506.
- [20]. Yehuda R, Engel SM, Brand SR, Seckl J, Marcus SM, Berkowitz GS. Transgenerational effects of posttraumatic stress disorder in babies of mothers exposed to the World Trade Center attacks during pregnancy. 2005;90(7):4115- 4118. *Journal of Clinical Endocrinology and Metabolism*.
- [21]. Khashan AS, Abel KM, McNamee R, Pedersen MG, Webb RT, Baker PN, Kenny LC, Mortensen PB. Higher risk of offspring schizophrenia following antenatal maternal exposure to severe adverse life events. *Archives of General Psychiatry*. 2008;65(2):146-152.
- [22]. Vivette Glover, MA, PhD, DSc . The Effects of Prenatal Stress on Child Behavioural and Cognitive Outcomes Start at the Beginning. Institute of Reproductive and Developmental Biology, Imperial College London, United Kingdom. *Encyclopedia on Early Childhood Development*. April 2019, Rev. ed.
- [23]. Scassellati C, Bonvicini C, Faraone SV, Gennarelli M. Biomarkers and attention-deficit/hyperactivity disorder: a systematic review and meta-analyses. *J Am Acad Child Adolesc Psychiatry*. 2012;51(10):1003-1019.e20.
- [24]. Franz AP, Bolat GU, Bolat H, Matijasevich A, Santos IS, Silveira RC, et al. Attention-deficit/hyperactivity disorder and very preterm/very low birth weight: a meta-analysis. *Pediatrics*. 2018;141(1). pii: e20171645.
- [25]. Langley K, Rice F, van den Bree MB, Thapar A. Maternal smoking during pregnancy as an environmental risk factor for attention deficit hyperactivity disorder behaviour. A review. *Minerva Pediatr*. 2005;57(6):359-71.
- [26]. Adeyemo BO, Biederman J, Zafonte R, Kagan E, Spencer TJ, Uchida M, et al. Mild traumatic brain injury and ADHD: a systematic review of the literature and meta-analysis. *J AttenDisord*. 2014;18(7):576-84.
- [27]. Stevens SE, Kumsta R, Kreppner JM, Brookes KJ, Rutter M, Sonuga-Barke EJ. Dopamine transporter gene polymorphism moderates the effects of severe deprivation on ADHD symptoms: developmental continuities in gene-environment interplay. *Am J Med Genet B Neuropsychiatr Genet*. 2009;150B(6):753-61.
- [28]. Bhutta AT, Cleves MA, Casey PH, Cradock MM, Anand KJ. Cognitive and behavioral outcomes of school-aged children who were born preterm: a meta-analysis. *JAMA*. 2002;288(6):728-37.
- [29]. Dong T, Hu W, Zhou X, Lin H, Lan L, Hang B, et al. Prenatal exposure to maternal smoking during pregnancy and attention-deficit/hyperactivity disorder in offspring: a meta-analysis. *ReprodToxicol*. 2018;76:63-70.
- [30]. Yoshimasu K, Kiyohara C, Takemura S, Nakai K. A meta-analysis of the evidence on the impact of prenatal and early infancy exposures to mercury on autism and attention deficit/hyperactivity disorder in the childhood. *Neurotoxicology*. 2014;44:121-31.
- [31]. Goodlad JK, Marcus DK, Fulton JJ. Lead and attention-deficit/hyperactivity disorder (ADHD) symptoms: a meta-analysis. *ClinPsychol Rev*. 2013;33(3):417-25.
- [32]. Khoshbakht Y, Bidaki R, Salehi-Abargouei A. Vitamin D status and attention deficit hyperactivity

- disorder: a systematic review and meta-analysis of observational studies. *AdvNutr*. 2018;9(1):9-20.
- [33]. Wolraich ML, Wilson DB, White JW. The effect of sugar on behavior or cognition in children. A meta-analysis. *JAMA*. 1995;274(20):1617-21.
- [34]. Thompson W, Russell G, Baragwanath G, Matthews J, Vaidya B, Thompson-Coon J. Maternal thyroid hormone insufficiency during pregnancy and risk of neurodevelopmental disorders in offspring: A systematic review and meta-analysis. *ClinEndocrinol (Oxf)*. 2018;88(4):575-584.
- [35]. Lundahl A, Kidwell KM, Van Dyk TR, Nelson TD. A meta-analysis of the effect of experimental sleep restriction on youth's attention and hyperactivity. *DevNeuropsychol*. 2015;40(3):104-21.
- [36]. Curran EA, O'Neill SM, Cryan JF, Kenny LC, Dinan TG, Khashan AS, et al. Research review: birth by caesarean section and development of autism spectrum disorder and attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *J Child Psychol Psychiatry*. 2015;56(5):500-8.
- [37]. Hoffmann MS, Polanczyk GV, Kieling C, Dos Santos IP, Willcutt EG, Rohde LA, et al. Attention-deficit/ hyperactivity disorder and solar irradiance: a cloudy perspective. *Biol Psychiatry*. 2014;76(8):e19-20.
- [38]. Hawkey E, Nigg JT. Omega-3 fatty acid and ADHD: blood level analysis and meta-analytic extension of supplementation trials. *ClinPsychol Rev* 2014;34(6):496–505. doi: 10.1016/j.cpr.2014.05.005.
- [39]. Rucklidge JJ, Frampton CM, Gorman B, Boggis A. Vitamin-mineral treatment of attention-deficit hyperactivity disorder in adults: double-blind randomised placebo-controlled trial. *Br J Psychiatry* 2014;204:306–15. doi: 10.1192/bjp.bp.113.132126.
- [40]. Nigg JT, Lewis K, Edinger T, Falk M. Meta-analysis of attention deficit/ hyperactivity disorder or attention-deficit/hyperactivity disorder symptoms, restriction diet, and synthetic food color additives. *J Am Acad Child Adolesc Psychiatry* 2012;51(1):86–97. e8.
- [41]. Heilskov Rytter MJ, Andersen LBB, Houmann T, Bilenberg N, Hvolby A, Molgaard C, Michaelsen KF, Lauritzen L. Diet in the treatment of ADHD in children—a systematic review of the literature. *Nord J Psychiatry* 2015;69(1):1–18.
- [42]. Eyles DW, Smith S, Kinobe R, Hewison M, McGrath JJ. Distribution of the vitamin D receptor and 1 α -hydroxylase in human brain. *J Chem Neuroanat* 2005;29(1):21–30.
- [43]. Cui X, Pertile R, Liu P, Eyles D. Vitamin D regulates tyrosine hydroxylase expression: N-cadherin a possible mediator. *Neuroscience* 2015;304:90–100.
- [44]. Patrick RP, Ames BN. Vitamin D and the omega-3 fatty acids control serotonin synthesis and action, part 2: relevance for ADHD, bipolar disorder, schizophrenia, and impulsive behavior. *FASEB J* 2015;29(6):2207–22. doi: 10.1096/fj.14-268342.
- [45]. Celik G, Taş D, Tahiroğlu A, Avci A, Yuksel B, Cam P. Vitamin D deficiency in obsessive-compulsive disorder patients with pediatric autoimmune neuropsychiatric disorders associated with streptococcal infections: a case control study. *NoroPsikiyatrisi* 2016;53(1):31–4. doi: 10.5152/npa.2015.8763.
- [46]. Sharif MR, Madani M, Tabatabaei F, Tabatabaee Z. The relationship between serum vitamin D level and attention deficit hyperactivity disorder. *Iran J Child Neurol* 2015;9(4):48–53.
- [47]. Kamal M, Bener A, Ehlayel MS. Is high prevalence of vitamin D deficiency a correlate for attention deficit hyperactivity disorder? *AttenDeficHyperactDisord* 2014;6(2):73–8. doi: 10.1007/s12402-014-

0130-5.

- [48]. Gustafsson P, Rylander L, Lindh CH, Jonsson BAG, Ode A, Olofsson P, Ivarsson SA, Rignell-Hydbom A, Haglund N, Kallen K. Vitamin D status at birth and future risk of attention deficit/hyperactivity disorder (ADHD). *PLoS One* 2015;10(10). doi: 10.1371/journal.pone.0140164.
- [49]. Shang-Guan LL, Zhao YR. [Serum levels of 25-hydroxyvitamin D in children with attention deficit hyperactivity disorder]. *Zhongguo Dang Dai ErKeZaZhi* 2015;17(8):837–40.
- [50]. Goksugur SB, Tufan AE, Semiz M, Gunes C, Bekdas M, Tosun M, Demircioglu F. Vitamin D status in children with attention deficit-hyperactivity disorder. *PediatrInt* 2014;56(4):515–9. doi: 10.1111/ped.12286.
- [51]. Bener A, Kamal M. Predict attention deficit hyperactivity disorder? Evidence-based medicine. *Glob J Health Sci* 2014;6(2):47–57. doi: 10.5539/gjhs.v6n2p47.
- [52]. Morales E, Julvez J, Torrent M, Ballester F, Rodriguez-Bernal CL, Andiarena A, Vegas O, Castilla AM, Rodriguez-Dehli C, Tardon A, et al. Vitamin D in pregnancy and attention deficit hyperactivity disorder-like symptoms in childhood. *Epidemiology* 2015;26(4):458–65. doi: 10.1097/EDE.0000000000000292.
- [53]. Downloaded from <https://academic.oup.com/advances/article-abstract/9/1/9/4848944> by guest on 30 July 2019
- [54]. Martel MM, Klump K, Nigg JT, Breedlove SM, Sisk CL. Potential hormonal mechanisms of attention-deficit/hyperactivity disorder and major depressive disorder: a new perspective. *HormBehav*. 2009;55(4):465–479.
- [55]. Davies W. Sex differences in attention deficit hyperactivity disorder: candidate genetic and endocrine mechanisms. *Front Neuroendocrinol*. 2014;35(3):331–346.
- [56]. Brunton PJ. Programming the brain and behaviour by early-life stress: a focus on neuroactive steroids. *J Neuroendocrinol*. 2015;27(6): 468–480.
- [57]. Hines M, Brook C, Conway GS: Androgen and psychosexual development: core gender identity, sexual orientation and recalled childhood gender role behavior in women and men with congenital adrenal hyperplasia (CAH). *J Sex Res* 2004;41:1-7
- [58]. Morris JA, Jordan CL, Breedlove SM. Sexual differentiation of the vertebrate nervous system. *Nat Neurosci*. 2004;7(10):1034–1039
- [59]. Baron –Cohen S, Lutchmaya S, Knickmeyer R: *Prenatal Testosterone in Mind* . Cambridge, MIT Press, 2004.
- [60]. Talge NM, Neal C, Glover V: Antenatal Maternal stress and long-term effects on child neurodevelopment: how and why? *J Child Psychol Psychiatry* 2007; 48:245-261.
- [61]. Mahone EM. Why fewer females than males are diagnosed with attention-deficit-hyperactivity disorder: it might not be hormones. *Dev Med Child Neurol*. 2010;52(9):790–791.
- [62]. Lemiere J, Boets B, Danckaerts M. No association between the 2D:4D fetal testosterone marker and multidimensional attentional abilities in children with ADHD. *Dev Med Child Neurol*. 2010;52(9):e202–e208.
- [63]. Liu J, Portnoy J, Raine A. Association between a marker for prenatal testosterone exposure and externalizing behavior problems in children. *DevPsychopathol*. 2012;24(3):771–782.

- [64]. Lee S. S. Humphreys K. L. Flory K. Liu R. Glass K. (2011). Prospective association of childhood attention-deficit/hyperactivity disorder (ADHD) and substance use and abuse/dependence: A meta-analytic review . *Clinical Psychology Review* , 31 , 328 – 341 . doi:10.1016/j.cpr.2011.01.006 [doi]
- [65]. Fuemmeler B. F. Kollins S. H. McClernon F. J. (2007). Attention deficit hyperactivity disorder symptoms predict nicotine dependence and progression to regular smoking from adolescence to young adulthood . *Journal of Pediatric Psychology* , 32 , 1203 – 1213 . doi:jsm051 [pii]
- [66]. Molina B. S. Pelham W. E. Jr. (2003). Childhood predictors of adolescent substance use in a longitudinal study of children with ADHD . *Journal of Abnormal Psychology* , 112 , 497 – 507 .
- [67]. Spencer T. J. Faraone S. V. Tarko L. McDermott K. Biederman J. (2014). Attention-deficit/hyperactivity disorder and adverse health outcomes in adults . *The Journal of Nervous and Mental Disease* , 202 , 725 – 731 .
- [68]. Bleck J. DeBate R. D. (2013). Exploring the co-morbidity of attention-deficit/hyperactivity disorder with eating disorders and disordered eating behaviors in a nationally representative community-based sample . *Eating Behaviors* , 14 , 390 – 393 . doi:10.1016/j.eatbeh.2013.05.009 [doi]
- [69]. Brown L. K. Hadley W. Stewart A. Lescano C. Whiteley L. Donenberg G. DiClemente R. (2010). Psychiatric disorders and sexual risk among adolescents in mental health treatment . *Journal of Consulting and Clinical Psychology* , 78 , 590 – 597 .
- [70]. Flory K. Molina B. S. Pelham W. E. Jr. Gnagy E. Smith B. (2006). Childhood ADHD predicts risky sexual behavior in young adulthood . *Journal of Clinical Child and Adolescent Psychology* , 35 , 571 – 577 . doi:10.1207/s15374424jccp3504_8 [doi]