



---

## **New Index for Salinity Assessment Applied on Saline Context Area (Case of the Lower Chélif Plain)**

Koulla Nouredine<sup>a\*</sup>, Mokhtari Djamel Eddine<sup>b</sup>, Douaoui Abd El Kader<sup>c</sup>

<sup>a</sup> *Institut of agronomic sciences, Hassiba Ben Bouali Chlef university. Es-Salem City, 19 national streets, 02000 Chlef, Algeria. Institut of agronomic sciences*

<sup>b</sup> *Laboratory of Ecology & Environment sciences, University of sciences & technology Houari-Boumedienne Algiers*

<sup>c</sup> *Laboratory of Agricultural production and sustainable development of natural resources. Khémis Miliana University*

### **Abstract**

The plain of the lower Chélif is located 220 km to the west of Algiers and extends over 60000 ha. This area is suffered by soil salinization in relation with various degradation levels. The remote-sensing data with salinity measured data of the ground surface has been used to evaluate the soil salinity using remote-sensing indices and apply a comparison the purpose index ISk. The correlation between the three applied indices have shown a good result in term of salinity assessment and made a better estimation of soil salinity.

**Keywords:** Salinity; indicator; remote sensing; Landsat TM; Lower Chélif.

### **1. Introduction**

Salinity is one of the most environmental issues affecting soil resources in the world [1, 2]. It causes severe environmental degradation that impedes crop growth and overall regional production [3, 4]. In Algeria, the salinity known an important increase at the last decade [5, 6], the lower Chélif is one of the most heavily regions affected by soil salinity problems in Algeria, with further of 80% are [5], due to irrigation activities [6].

---

\* Corresponding author. Tel.: +213772222295.

E-mail address: koulla-033@hotmail.com.

Several indicators were used for the estimation of salinity by the use of remote sensing in the world [6,7,8,9].

The proposed salinity index (ISk Koulla) is applicable to make a good assessment based on acquire data in the visible and near-infrared satellite imagery bands. They have been used in our study area, especially in the context of salinity.

## **2. Materials and methods**

### ***2.1 Study area***

The Lower Chéiff Plain, which encompasses over 60,000 ha in land area; it is located in the extreme northern part of the Chéiff Basin in Northwest Algeria, between longitudes 0°40 and 01°06'08"E and latitudes 34°03'12"and 36°05'57"N. The Lower Chéiff specific climate is of semi-arid with very hot summers, low winter temperatures and a mean annual rainfall of 250 mm [6].

### ***2.2. Satellite image data***

The choice of the image type referred from a satellite to another, For our study we used the raw image Landsat TM acquired in June 2006.

### ***2.3. The soil sampling***

The ground recognition and the sample selection test were carried out during one period being spread out over July 2006. Each sampling site was georeferenced and located on the ground by the use of a GPS into UTM projection (Zone 31 North, WGS 84). The full number of the 400 taken samples is distributed on all the plain.

### ***2.4. The laboratory Analyzes***

The analyzes at the laboratory were based on the measure of electric conductivity using diluted focal (1/5) for all 400 taken samples of the surface layer (0 - 20cm).

### ***2.5. Salinity used indices***

Three salinity indices were established using the same landsat TM satellite image (table 1)

## **3. Results and Discussions**

In general, the three indices IS, NDSI, and ISK that were extracted from the Landsat satellite image were used for assessing the soil salinity for all 400 ground-measured samples.

The ISK indicator give a good correlation values 0,70 ; 0,75 respectively with IS, NDSI index. [6, 8]. (Table. 2). However, the results of Bannari and his colleagues [10], indicated that the NDSI index was a poor indicator for salt concentrations [4].

**Table 1.** Summary of soil salinity indices used

Index	Formulation	Reference
Normalized Differential Salinity Index (NDSI)	$NDSI = (R - NIR) / (R + NIR)$	[8]
Salinity Index (IS)	$IS = \sqrt{G^2 + R^2 + NIR^2}$	[6]
Salinity Index (ISK)	$ISK = \frac{\sqrt{(R - G)(R + G)}}{\sqrt{R^2 + G^2}}$	The proposed salinity index (IS Koulla)

**Table 2.** Correlation between ISK and other soil salinity used indices

Index	NDSI	IS	ISK
Normalized Differential Salinity Index (NDSI)	1,00	0,63	0,75
Salinity Index (IS)	0,63	1,00	0,70
Salinity Index (ISK Koulla)	0,75	0,70	1,00

#### 4. Conclusion

Overall, the IS, NDSI and ISk indices were useful for assessing the soil salinity. This study indicated that The Isk can be considered as a new index for salinity assessment, particularly in the context of salinity.

#### References

- [1] Fernandez-Buces N., Siebe C., Cramb. S, Palacio J. 2006. Mapping soil salinity using a com-bined spectral response index for bare soil and vegetation: A case study in the former lake Texcoco, Mexico. Journal of Arid Environments 65 (2006) 644– 667.
- [2] Metternicht,G., Zinck,J.A., 2008. Remote Sensing of Soil Salinization: Impact on Land Management.CRC Press, Taylor and Francis, NewYork.
- [3] Akhtar Abbas , Shahbaz Khan, Nisar Hussain, Munir Hanjra, Saud Akbar., 2013. Characterizing soil salinity in irrigated agriculture using a remote sensing approach Physicsand Chemistry of the Earth. 55-57, 43–52.

- [4] Amal Allbed, Lalit Kumar, Yousef Y. Aldakheel., 2014. Assessing soil salinity using soil salinity and vegetation indices derived From IKONOS high-spatial resolution imageries: Applications in a date palm dominated region. *Geoderma* 230–231. 1-8
- [5] Douaoui A ., 2005. Variabilité Spatiale de la salinité et sa relation avec certaines caractéristiques des sols de la plaine du Bas-Chélif .Apport de la géostatistique et de la télédétection .Thèse Doctorat d'état, INA Alger, 233p.
- [6] Douaoui, A.E.K .,Nicolas,H .,Walter,C .,2006. Detecting salinity hazards with in a semiarid Context by means of combining soil and remote-sensing data. *Geoderma*134 (1), 217–230.
- [7] Tripathi, N. K ., Brijesh , K . R .,D ., P., 1997. Spatial modelling of soil alkalinity in GIS environment using IRS data. Paper presented at the18 th Asian Conference in Remote Sensing.
- [8] Khan, N.M., Rastoskuev,V.V., Sato,Y., Shiozawa,S., 2005. Assessment of hydro saline land degradation by using a simple approach of remote sensing indicators. *Agric. Water Manag.* 77(1),96–109.
- [9] Abbas,A.,Khan,S.,2007. Using remote sensing techniques for appraisal of irrigated soil salinity. Paper presented at the Advances and Applications for Management and Decision Making Land, Water and Environmental Management: Integrated Systems For Sustainability MODSIM07.
- [10] Bannari,A., Guedon ,A.M .,El-Harti ,A .,Cherkaoui,F.Z., El-Ghmari,A., 2008. Characterization of slightly and moderately saline and sodic soils in irrigated agricultural land using simulated data of advanced land imaging (EO-1) sensor .*Commun. Soil Sci. Plant Anal.* 39(19–20), 2795–2811.